# **NCERT Solutions for Class 11 Maths Maths Chapter 4**

## **Principle of Mathematical Induction Class 11**

Chapter 4 Principle of Mathematical Induction Exercise 4.1 Solutions

Exercise 4.1 : Solutions of Questions on Page Number : 94 Q1 :

Prove the following by using the principle of mathematical induction for all  $n \in N$ :

$$1 + 3 + 32 + \dots + 3n-1 = \frac{(3^{n} - 1)}{2}$$

#### Answer :

Let the given statement be P(n), i.e.,

$$\frac{\left(3^n-1\right)}{2}$$

$$P(n): 1 + 3 + 3^{2} + \dots + 3^{n \hat{a} \in 1} = 2$$

For n = 1, we have

P(1): 1 = 
$$\frac{(3^{1}-1)}{2} = \frac{3-1}{2} = \frac{2}{2} = 1$$
, which is true.

Let P(k) be true for some positive integer k, i.e.,

$$1 + 3 + 32 + ... + 3k-1 = \frac{(3k - 1)}{2} \qquad ...(i)$$

We shall now prove that P(k + 1) is true.

#### Consider

 $1 + 3 + 3^{2} + \dots + 3^{k a \in 1} + 3^{(k+1) a \in 1}$  $= (1 + 3 + 3^{2} + \dots + 3^{k a \in 1}) + 3^{k}$ 

$$= \frac{(3^{k} - 1)}{2} + 3^{k} \qquad [Using (i)]$$
$$= \frac{(3^{k} - 1) + 2 \cdot 3^{k}}{2}$$
$$= \frac{(1 + 2) 3^{k} - 1}{2}$$
$$= \frac{3 \cdot 3^{k} - 1}{2}$$
$$= \frac{3^{k+1} - 1}{2}$$

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

## Q2 :

Prove the following by using the principle of mathematical induction for

$$1^{3} + 2^{3} + 3^{3} + \dots + n^{3} = \left(\frac{n(n+1)}{2}\right)^{2}$$

all  $n \in N$ :

#### Answer :

Let the given statement be P(n), i.e.,

P(n): 
$$1^3 + 2^3 + 3^3 + \dots + n^3 = \left(\frac{n(n+1)}{2}\right)^2$$

For n = 1, we have

P(1): 1<sup>3</sup> = 1 = 
$$\left(\frac{1(1+1)}{2}\right)^2 = \left(\frac{1.2}{2}\right)^2 = 1^2 = 1$$
, which is true

Let P(k) be true for some positive integer k, i.e.,

$$1^{3} + 2^{3} + 3^{3} + \dots + k^{3} = \left(\frac{k(k+1)}{2}\right)^{2} \qquad \dots (i)$$

We shall now prove that P(k + 1) is true.

Consider

 $1^3 + 2^3 + 3^3 + \dots + k^3 + (k+1)^3$ 

$$= \left(\frac{k(k+1)}{2}\right)^{2} + (k+1)^{3} \qquad [Using (i)]$$

$$= \frac{k^{2}(k+1)^{2}}{4} + (k+1)^{3}$$

$$= \frac{k^{2}(k+1)^{2} + 4(k+1)^{3}}{4}$$

$$= \frac{(k+1)^{2} \left\{k^{2} + 4(k+1)\right\}}{4}$$

$$= \frac{(k+1)^{2} \left\{k^{2} + 4k + 4\right\}}{4}$$

$$= \frac{(k+1)^{2} (k+2)^{2}}{4}$$

$$= \frac{(k+1)^{2} (k+1+1)^{2}}{4}$$

$$= \left(\frac{(k+1)(k+1+1)}{2}\right)^{2}$$

 $=(1^{3}+2^{3}+3)$ 

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

## Q3 :

Prove the following by using the principle of mathematical induction for

all 
$$n \in N$$
:  $1 + \frac{1}{(1+2)} + \frac{1}{(1+2+3)} + \dots + \frac{1}{(1+2+3+\dots n)} = \frac{2n}{(n+1)}$ 

#### Answer :

Let the given statement be P(n), i.e.,

P(n): 
$$1 + \frac{1}{1+2} + \frac{1}{1+2+3} + \dots + \frac{1}{1+2+3+\dots n} = \frac{2n}{n+1}$$

For n = 1, we have

P(1): 1 = 
$$\frac{2.1}{1+1} = \frac{2}{2} = 1$$
 which is true.

Let P(k) be true for some positive integer k, i.e.,

$$1 + \frac{1}{1+2} + \dots + \frac{1}{1+2+3} + \dots + \frac{1}{1+2+3+\dots+k} = \frac{2k}{k+1} \qquad \dots (i)$$

We shall now prove that P(k + 1) is true.

Consider

$$\begin{aligned} 1 + \frac{1}{1+2} + \frac{1}{1+2+3} + \dots + \frac{1}{1+2+3+\dots+k} + \frac{1}{1+2+3+\dots+k+(k+1)} \\ &= \left(1 + \frac{1}{1+2} + \frac{1}{1+2+3} + \dots + \frac{1}{1+2+3+\dots+k}\right) + \frac{1}{1+2+3+\dots+k+(k+1)} \\ &= \frac{2k}{k+1} + \frac{1}{1+2+3+\dots+k+(k+1)} \\ &= \frac{2k}{k+1} + \frac{1}{\left(\frac{(k+1)(k+1+1)}{2}\right)} \\ &= \frac{2k}{(k+1)} + \frac{2}{(k+1)(k+2)} \\ &= \frac{2}{(k+1)} \left(k + \frac{1}{k+2}\right) \\ &= \frac{2}{(k+1)} \left(\frac{k(k+2)+1}{k+2}\right) \\ &= \frac{2}{(k+1)} \left(\frac{k^2+2k+1}{k+2}\right) \\ &= \frac{2(k+1)}{(k+1)(k+2)} \\ &= \frac{2(k+1)}{(k+1)(k+2)} \end{aligned}$$

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

Q4 :

Prove the following by using the principle of mathematical induction for all  $n \in N$ : 1.2.3 + 2.3.4 + ... + n(n + 1)

$$\frac{n(n+1)(n+2)(n+3)}{4}$$

(n+2) = 4

#### Answer :

Let the given statement be P(n), i.e.,

$$P(n): 1.2.3 + 2.3.4 + \dots + n(n+1)(n+2) = \frac{n(n+1)(n+2)(n+3)}{4}$$

For n = 1, we have

P(1): 1.2.3 = 6 = 
$$\frac{1(1+1)(1+2)(1+3)}{4} = \frac{1.2.3.4}{4} = 6$$
, which is true.

Let P(k) be true for some positive integer k, i.e.,

$$\frac{k(k+1)(k+2)(k+3)}{4} \dots (i)$$

We shall now prove that P(k + 1) is true.

Consider

$$\begin{split} 1.2.3 + 2.3.4 + \ldots + k(k+1) & (k+2) + (k+1) & (k+2) & (k+3) \\ = & \{1.2.3 + 2.3.4 + \ldots + k(k+1) & (k+2)\} + (k+1) & (k+2) & (k+3) \end{split}$$

$$= \frac{k(k+1)(k+2)(k+3)}{4} + (k+1)(k+2)(k+3) \qquad [Using (i)]$$
  
=  $(k+1)(k+2)(k+3)\left(\frac{k}{4}+1\right)$   
=  $\frac{(k+1)(k+2)(k+3)(k+4)}{4}$   
=  $\frac{(k+1)(k+1+1)(k+1+2)(k+1+3)}{4}$ 

Thus, P(k + 1) is true whenever P(k) is

true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

## Q5 :

#### Prove the following by using the principle of mathematical induction for

$$1.3 + 2.3^{2} + 3.3^{3} + \dots + n.3^{n} = \frac{(2n-1)3^{n+1} + 3}{4}$$

all  $n \in N$ 

#### Answer :

Let the given statement be P(n), i.e.,

P(n): 
$$1.3 + 2.3^{2} + 3.3^{3} + \dots + n3^{n} = \frac{(2n-1)3^{n+1} + 3}{4}$$

For n = 1, we have

P(1): 1.3 = 3 = 
$$\frac{(2.1-1)3^{1+1}+3}{4} = \frac{3^2+3}{4} = \frac{12}{4} = 3$$
, which is true.

Let P(k) be true for some positive integer k, i.e.,

$$1.3 + 2.3^{2} + 3.3^{3} + \dots + k3^{k} = \frac{(2k-1)3^{k+1} + 3}{4} \qquad \dots (i)$$

We shall now prove that P(k + 1) is true.

### Consider

$$1.3 + 2.3^{2} + 3.3^{3} + ... + k3^{k} + (k + 1) 3^{k+1}$$

$$= (1.3 + 2.3^{2} + 3.3^{3} + ... + k3^{k}) + (k + 1) 3^{k+1}$$

$$= \frac{(2k - 1) 3^{k+1} + 3}{4} + (k + 1) 3^{k+1}}{4}$$

$$= \frac{(2k - 1) 3^{k+1} + 3 + 4(k + 1) 3^{k+1}}{4}$$

$$= \frac{3^{k+1} \{2k - 1 + 4(k + 1)\} + 3}{4}$$

$$= \frac{3^{k+1} \{6k + 3\} + 3}{4}$$

$$= \frac{3^{k+1} \{6k + 3\} + 3}{4}$$

$$= \frac{3^{(k+1)+1} \{2k + 1\} + 3}{4}$$

$$= \frac{3^{(k+1)+1} \{2k + 1\} + 3}{4}$$

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

## Q6:

Prove the following by using the principle of mathematical induction for

$$1.2 + 2.3 + 3.4 + \dots + n.(n+1) = \left[\frac{n(n+1)(n+2)}{3}\right]$$

all  $n \in N$ :

#### Answer :

Let the given statement be P(n), i.e.,

P(n):  

$$1.2 + 2.3 + 3.4 + \dots + n.(n+1) = \left[\frac{n(n+1)(n+2)}{3}\right]$$

For n = 1, we have

P(1): 
$$1.2 = 2 = \frac{1(1+1)(1+2)}{3} = \frac{1.2.3}{3} = 2$$
, which is true.

Let P(k) be true for some positive integer k, i.e.,

$$1.2 + 2.3 + 3.4 + \dots + k.(k+1) = \left[\frac{k(k+1)(k+2)}{3}\right] \dots (i)$$

We shall now prove that P(k + 1) is true.

Consider

$$1.2 + 2.3 + 3.4 + \dots + k.(k + 1) + (k + 1).(k + 2)$$
  
=  $[1.2 + 2.3 + 3.4 + \dots + k.(k + 1)] + (k + 1).(k + 2)$   
=  $\frac{k(k+1)(k+2)}{3} + (k+1)(k+2)$  [Using (i)]  
=  $(k+1)(k+2)(\frac{k}{3}+1)$   
=  $\frac{(k+1)(k+2)(k+3)}{3}$   
=  $\frac{(k+1)(k+1+1)(k+1+2)}{3}$ 

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

#### Q7 :

Prove the following by using the principle of mathematical induction for

$$1.3+3.5+5.7+...+(2n-1)(2n+1) = \frac{n(4n^2+6n-1)}{3}$$

all  $n \in N$ :

#### Answer :

Let the given statement be P(n), i.e.,

P(n): 
$$1.3 + 3.5 + 5.7 + ... + (2n-1)(2n+1) = \frac{n(4n^2 + 6n-1)}{3}$$

For n = 1, we have

P(1):1.3 = 3 = 
$$\frac{1(4.1^2 + 6.1 - 1)}{3} = \frac{4 + 6 - 1}{3} = \frac{9}{3} = 3$$
, which is true.

Let P(k) be true for some positive integer k, i.e.,

$$1.3 + 3.5 + 5.7 + \dots + (2k-1)(2k+1) = \frac{k(4k^2 + 6k - 1)}{3} \dots (i)$$

We shall now prove that P(k + 1) is true.

$$(1.3 + 3.5 + 5.7 + ... + (2k a \in "1) (2k + 1) + {2(k + 1) a \in "1} {2(k + 1) + 1}$$

$$= \frac{k(4k^{2} + 6k - 1)}{3} + (2k + 2 - 1)(2k + 2 + 1) \qquad [Using (i)]$$

$$= \frac{k(4k^{2} + 6k - 1)}{3} + (2k + 1)(2k + 3)$$

$$= \frac{k(4k^{2} + 6k - 1) + 3(4k^{2} + 8k + 3)}{3}$$

$$= \frac{4k^{3} + 6k^{2} - k + 12k^{2} + 24k + 9}{3}$$

$$= \frac{4k^{3} + 18k^{2} + 23k + 9}{3}$$

$$= \frac{4k^{3} + 14k^{2} + 9k + 4k^{2} + 14k + 9}{3}$$

$$= \frac{k(4k^{2} + 14k + 9) + 1(4k^{2} + 14k + 9)}{3}$$

$$=\frac{(k+1)\left\{4k^{2}+8k+4+6k+6-1\right\}}{3}$$
$$=\frac{(k+1)\left\{4(k^{2}+2k+1)+6(k+1)-1\right\}}{3}$$
$$=\frac{(k+1)\left\{4(k+1)^{2}+6(k+1)-1\right\}}{3}$$

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

#### Q8 :

Prove the following by using the principle of mathematical induction for all  $n \in N$ :  $1.2 + 2.2^2 + 3.2^2 + ... + n.2^n = (n - 1) 2^{n+1} + 2$ 

#### Answer :

Let the given statement be P(n), i.e.,

P(*n*): 1.2 + 2.2<sup>2</sup> + 3.2<sup>2</sup> + ... + *n*.2<sup>*n*</sup> = (*n* – 1) 2<sup>*n*+1</sup> + 2

For n = 1, we have

P(1): 1.2 = 2 = (1 – 1)  $2^{1+1}$  + 2 = 0 + 2 = 2, which is true.

Let P(k) be true for some positive integer k, i.e.,

 $1.2 + 2.2^2 + 3.2^2 + ... + k.2^k = (k \hat{a} \in 1) 2^{k+1} + 2 ... (i)$ 

We shall now prove that P(k + 1) is true.

Consider

$$\{1.2 + 2.2^{2} + 3.2^{3} + \dots + k.2^{k}\} + (k+1) \cdot 2^{k+1}$$
  
=  $(k-1)2^{k+1} + 2 + (k+1)2^{k+1}$   
=  $2^{k+1}\{(k-1) + (k+1)\} + 2$   
=  $2^{k+1}.2k + 2$   
=  $k.2^{(k+1)+1} + 2$   
=  $\{(k+1)-1\}2^{(k+1)+1} + 2$ 

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

Prove the following by using the principle of mathematical induction for

all 
$$n \in N$$
:  $\frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots + \frac{1}{2^n} = 1 - \frac{1}{2^n}$ 

### Answer :

Let the given statement be P(n), i.e.,

P(n): 
$$\frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots + \frac{1}{2^n} = 1 - \frac{1}{2^n}$$

For n = 1, we have

P(1): 
$$\frac{1}{2} = 1 - \frac{1}{2^1} = \frac{1}{2}$$
, which is true

Let P(k) be true for some positive integer k, i.e.,

$$\frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots + \frac{1}{2^{k}} = 1 - \frac{1}{2^{k}} \qquad \dots (i)$$

We shall now prove that P(k + 1) is true.

## Consider

$$\begin{pmatrix} \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots + \frac{1}{2^{k}} \end{pmatrix} + \frac{1}{2^{k+1}}$$

$$= \left(1 - \frac{1}{2^{k}}\right) + \frac{1}{2^{k+1}}$$

$$= 1 - \frac{1}{2^{k}} + \frac{1}{2 \cdot 2^{k}}$$

$$= 1 - \frac{1}{2^{k}} \left(1 - \frac{1}{2}\right)$$

$$= 1 - \frac{1}{2^{k}} \left(\frac{1}{2}\right)$$

$$= 1 - \frac{1}{2^{k+1}}$$

$$[Using (i)]$$

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

## Q10:

Prove the following by using the principle of mathematical induction for

all 
$$n \in N$$
:  
 $\frac{1}{2.5} + \frac{1}{5.8} + \frac{1}{8.11} + \dots + \frac{1}{(3n-1)(3n+2)} = \frac{n}{(6n+4)}$ 

## Answer :

Let the given statement be P(n), i.e.,

$$\frac{1}{2.5} + \frac{1}{5.8} + \frac{1}{8.11} + \dots + \frac{1}{(3n-1)(3n+2)} = \frac{n}{(6n+4)}$$

For n = 1, we have

$$P(1) = \frac{1}{2.5} = \frac{1}{10} = \frac{1}{6.1+4} = \frac{1}{10}$$
, which is true

Let P(k) be true for some positive integer k, i.e.,

$$\frac{1}{2.5} + \frac{1}{5.8} + \frac{1}{8.11} + \dots + \frac{1}{(3k-1)(3k+2)} = \frac{k}{6k+4} \qquad \dots (i)$$

We shall now prove that P(k + 1) is true.

$$\frac{1}{2.5} + \frac{1}{5.8} + \frac{1}{8.11} + \dots + \frac{1}{(3k-1)(3k+2)} + \frac{1}{\{3(k+1)-1\}} \{3(k+1)+2\}$$

$$= \frac{k}{6k+4} + \frac{1}{(3k+3-1)(3k+3+2)} \qquad [Using (i)]$$

$$= \frac{k}{6k+4} + \frac{1}{(3k+2)(3k+5)}$$

$$= \frac{k}{2(3k+2)} + \frac{1}{(3k+2)(3k+5)}$$

$$= \frac{1}{(3k+2)} \left(\frac{k}{2} + \frac{1}{3k+5}\right)$$

$$= \frac{1}{(3k+2)} \left(\frac{k(3k+5)+2}{2(3k+5)}\right)$$

$$= \frac{1}{(3k+2)} \left(\frac{3k^2+5k+2}{2(3k+5)}\right)$$

$$= \frac{1}{(3k+2)} \left(\frac{(3k+2)(k+1)}{2(3k+5)}\right)$$

$$= \frac{(k+1)}{6k+10}$$

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

## Q11 :

Prove the following by using the principle of mathematical induction for

all 
$$n \in N$$
:  
 $\frac{1}{1.2.3} + \frac{1}{2.3.4} + \frac{1}{3.4.5} + \dots + \frac{1}{n(n+1)(n+2)} = \frac{n(n+3)}{4(n+1)(n+2)}$ 

## Answer :

Let the given statement be P(n), i.e.,

$$\frac{1}{1.2.3} + \frac{1}{2.3.4} + \frac{1}{3.4.5} + \dots + \frac{1}{n(n+1)(n+2)} = \frac{n(n+3)}{4(n+1)(n+2)}$$

For n = 1, we have

$$P(1):\frac{1}{1\cdot 2\cdot 3} = \frac{1\cdot (1+3)}{4(1+1)(1+2)} = \frac{1\cdot 4}{4\cdot 2\cdot 3} = \frac{1}{1\cdot 2\cdot 3}, \text{ which is true.}$$

Let P(k) be true for some positive integer k, i.e.,

$$\frac{1}{1\cdot 2\cdot 3} + \frac{1}{2\cdot 3\cdot 4} + \frac{1}{3\cdot 4\cdot 5} + \dots + \frac{1}{k(k+1)(k+2)} = \frac{k(k+3)}{4(k+1)(k+2)} \qquad \dots (i)$$

We shall now prove that P(k + 1) is true.

$$\begin{bmatrix} \frac{1}{1\cdot 2\cdot 3} + \frac{1}{2\cdot 3\cdot 4} + \frac{1}{3\cdot 4\cdot 5} + \dots + \frac{1}{k(k+1)(k+2)} \end{bmatrix} + \frac{1}{(k+1)(k+2)(k+3)}$$

$$= \frac{k(k+3)}{4(k+1)(k+2)} + \frac{1}{(k+1)(k+2)(k+3)} \qquad [Using (i)]$$

$$= \frac{1}{(k+1)(k+2)} \left\{ \frac{k(k+3)^2 + 4}{4(k+3)} \right\}$$

$$= \frac{1}{(k+1)(k+2)} \left\{ \frac{k(k+3)^2 + 4}{4(k+3)} \right\}$$

$$= \frac{1}{(k+1)(k+2)} \left\{ \frac{k^3 + 6k^2 + 9k + 4}{4(k+3)} \right\}$$

$$= \frac{1}{(k+1)(k+2)} \left\{ \frac{k^3 + 2k^2 + k + 4k^2 + 8k + 4}{4(k+3)} \right\}$$

$$= \frac{1}{(k+1)(k+2)} \left\{ \frac{k(k+1)^2 + 4(k+1)^2}{4(k+3)} \right\}$$

$$= \frac{1}{(k+1)(k+2)} \left\{ \frac{k(k+1)^2 + 4(k+1)^2}{4(k+3)} \right\}$$

$$= \frac{(k+1)^2(k+4)}{4(k+1)(k+2)(k+3)}$$

$$= \frac{(k+1)^2(k+4)}{4(k+1)+1} \left\{ (k+1) + 2 \right\}$$

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

## Q12 :

Prove the following by using the principle of mathematical induction for

$$a + ar + ar^{2} + \dots + ar^{n-1} = \frac{a(r^{n} - 1)}{r - 1}$$
  
all  $n \in N$ :

Answer :

Let the given statement be P(n), i.e.,

$$P(n): a + ar + ar^{2} + ... + ar^{n-1} = \frac{a(r^{n}-1)}{r-1}$$

For n = 1, we have

$$P(1): a = \frac{a(r^1 - 1)}{(r - 1)} = a$$
, which is true.

Let P(k) be true for some positive integer k, i.e.,

$$a + ar + ar^{2} + \dots + ar^{k-1} = \frac{a(r^{k} - 1)}{r - 1} \dots$$
(i)

We shall now prove that P(k + 1) is true.

Consider

$$\{a + ar + ar^{2} + \dots + ar^{k-1}\} + ar^{(k+1)-1}$$

$$= \frac{a(r^{k} - 1)}{r - 1} + ar^{k} \qquad [Using(i)]$$

$$= \frac{a(r^{k} - 1) + ar^{k} (r - 1)}{r - 1}$$

$$= \frac{a(r^{k} - 1) + ar^{k+1} - ar^{k}}{r - 1}$$

$$= \frac{ar^{k} - a + ar^{k+1} - ar^{k}}{r - 1}$$

$$= \frac{ar^{k+1} - a}{r - 1}$$

$$= \frac{a(r^{k+1} - 1)}{r - 1}$$

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

## Q13:

## Prove the following by using the principle of mathematical induction for

$$\| n \in N: \left( 1 + \frac{3}{1} \right) \left( 1 + \frac{5}{4} \right) \left( 1 + \frac{7}{9} \right) \dots \left( 1 + \frac{(2n+1)}{n^2} \right) = \left( n+1 \right)^2$$

al

Let the given statement be P(n), i.e.,

$$P(n): \left(1+\frac{3}{1}\right) \left(1+\frac{5}{4}\right) \left(1+\frac{7}{9}\right) \dots \left(1+\frac{(2n+1)}{n^2}\right) = (n+1)^2$$

For n = 1, we have

P(1): 
$$\left(1+\frac{3}{1}\right) = 4 = \left(1+1\right)^2 = 2^2 = 4$$
, which is true.

Let P(k) be true for some positive integer k, i.e.,

$$\left(1+\frac{3}{1}\right)\left(1+\frac{5}{4}\right)\left(1+\frac{7}{9}\right)\dots\left(1+\frac{(2k+1)}{k^2}\right) = (k+1)^2 \qquad \dots (1)$$

We shall now prove that P(k + 1) is true.

Consider

$$\begin{bmatrix} \left(1+\frac{3}{1}\right)\left(1+\frac{5}{4}\right)\left(1+\frac{7}{9}\right)...\left(1+\frac{(2k+1)}{k^2}\right) \end{bmatrix} \left\{1+\frac{\left\{2(k+1)+1\right\}}{(k+1)^2}\right\}$$
  
=  $(k+1)^2 \left(1+\frac{2(k+1)+1}{(k+1)^2}\right)$  [Using(1)]  
=  $(k+1)^2 \left[\frac{(k+1)^2+2(k+1)+1}{(k+1)^2}\right]$   
=  $(k+1)^2+2(k+1)+1$   
=  $\{(k+1)+1\}^2$ 

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

## Q14 :

## Prove the following by using the principle of mathematical induction for

all 
$$n \in N$$
:  $\left(1 + \frac{1}{1}\right)\left(1 + \frac{1}{2}\right)\left(1 + \frac{1}{3}\right)...\left(1 + \frac{1}{n}\right) = (n+1)$ 

Answer :

Let the given statement be P(n), i.e.,

$$P(n):\left(1+\frac{1}{1}\right)\left(1+\frac{1}{2}\right)\left(1+\frac{1}{3}\right)...\left(1+\frac{1}{n}\right)=(n+1)$$

For n = 1, we have

$$P(1):(1+\frac{1}{1})=2=(1+1)$$
, which is true.

Let P(k) be true for some positive integer k, i.e.,

$$P(k):\left(1+\frac{1}{1}\right)\left(1+\frac{1}{2}\right)\left(1+\frac{1}{3}\right)...\left(1+\frac{1}{k}\right)=(k+1) \qquad ... (1)$$

We shall now prove that P(k + 1) is true.

Consider

$$\begin{bmatrix} \left(1+\frac{1}{1}\right)\left(1+\frac{1}{2}\right)\left(1+\frac{1}{3}\right)...\left(1+\frac{1}{k}\right)\end{bmatrix}\left(1+\frac{1}{k+1}\right) \\ = (k+1)\left(1+\frac{1}{k+1}\right) \\ = (k+1)\left(\frac{(k+1)+1}{(k+1)}\right) \\ = (k+1)+1 \end{bmatrix}$$
[Using (1)]

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

## Q15 :

## Prove the following by using the principle of mathematical induction for

$$1^{2} + 3^{2} + 5^{2} + \dots + (2n-1)^{2} = \frac{n(2n-1)(2n+1)}{3}$$

all  $n \in N$ :

## Answer :

Let the given statement be P(n), i.e.,

$$P(n) = 1^{2} + 3^{2} + 5^{2} + ... + (2n-1)^{2} = \frac{n(2n-1)(2n+1)}{3}$$

For n = 1, we have

$$P(1) = 1^2 = 1 = \frac{1(2.1-1)(2.1+1)}{3} = \frac{1.1.3}{3} = 1$$
, which is true.

Let P(k) be true for some positive integer k, i.e.,

$$P(k) = 1^{2} + 3^{2} + 5^{2} + \dots + (2k-1)^{2} = \frac{k(2k-1)(2k+1)}{3} \qquad \dots (1)$$

We shall now prove that P(k + 1) is true.

Consider

$$\{l^{2} + 3^{2} + 5^{2} + ... + (2k-1)^{2}\} + \{2(k+1)-1\}^{2}$$

$$= \frac{k(2k-1)(2k+1)}{3} + (2k+2-1)^{2}$$

$$[Using (1)]$$

$$= \frac{k(2k-1)(2k+1) + 3(2k+1)^{2}}{3}$$

$$= \frac{(2k+1)\{k(2k-1)+3(2k+1)\}}{3}$$

$$= \frac{(2k+1)\{2k^{2}-k+6k+3\}}{3}$$

$$= \frac{(2k+1)\{2k^{2}+5k+3\}}{3}$$

$$= \frac{(2k+1)\{2k^{2}+2k+3k+3\}}{3}$$

$$= \frac{(2k+1)\{2k(k+1)+3(k+1)\}}{3}$$

$$= \frac{(2k+1)\{2k(k+1)+3(k+1)\}}{3}$$

$$= \frac{(2k+1)\{2(k+1)-1\}\{2(k+1)+1\}}{3}$$

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

## Q16 :

Prove the following by using the principle of mathematical induction for

all 
$$n \in N$$
:  
 $\frac{1}{1.4} + \frac{1}{4.7} + \frac{1}{7.10} + \dots + \frac{1}{(3n-2)(3n+1)} = \frac{n}{(3n+1)}$ 

#### Answer :

Let the given statement be P(n), i.e.,

$$P(n): \frac{1}{1.4} + \frac{1}{4.7} + \frac{1}{7.10} + \dots + \frac{1}{(3n-2)(3n+1)} = \frac{n}{(3n+1)}$$

For n = 1, we have

$$P(1) = \frac{1}{1.4} = \frac{1}{3.1+1} = \frac{1}{4} = \frac{1}{1.4}$$
, which is true.

Let P(k) be true for some positive integer k, i.e.,

$$P(k) = \frac{1}{1.4} + \frac{1}{4.7} + \frac{1}{7.10} + \dots + \frac{1}{(3k-2)(3k+1)} = \frac{k}{3k+1} \qquad \dots (1)$$

We shall now prove that P(k + 1) is true.

Consider

1

$$\begin{cases} \frac{1}{1.4} + \frac{1}{4.7} + \frac{1}{7.10} + \dots + \frac{1}{(3k-2)(3k+1)} \\ + \frac{1}{\{3(k+1)-2\}\{3(k+1)+1\}} \\ = \frac{k}{3k+1} + \frac{1}{(3k+1)(3k+4)} \\ = \frac{1}{(3k+1)} \\ \left\{ k + \frac{1}{(3k+4)} \\ \\ = \frac{1}{(3k+1)} \\ \left\{ \frac{k(3k+4)+1}{(3k+4)} \\ \\ \\ = \frac{1}{(3k+1)} \\ \left\{ \frac{3k^2 + 4k + 1}{(3k+4)} \\ \\ \\ \\ = \frac{1}{(3k+1)} \\ \left\{ \frac{3k^2 + 3k + k + 1}{(3k+4)} \\ \\ \\ \\ \\ \\ = \frac{(3k+1)(k+1)}{(3k+4)} \\ \\ \\ \\ \\ = \frac{(k+1)}{3(k+1)+1} \end{cases}$$

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

Prove the following by using the principle of mathematical induction for

all 
$$n \in N$$
:  
 $\frac{1}{3.5} + \frac{1}{5.7} + \frac{1}{7.9} + \dots + \frac{1}{(2n+1)(2n+3)} = \frac{n}{3(2n+3)}$ 

## Answer :

Let the given statement be P(n), i.e.,

$$P(n):\frac{1}{3.5} + \frac{1}{5.7} + \frac{1}{7.9} + \dots + \frac{1}{(2n+1)(2n+3)} = \frac{n}{3(2n+3)}$$

For n = 1, we have

$$P(1): \frac{1}{3.5} = \frac{1}{3(2.1+3)} = \frac{1}{3.5}$$
, which is true.

Let P(k) be true for some positive integer k, i.e.,

$$P(k): \frac{1}{3.5} + \frac{1}{5.7} + \frac{1}{7.9} + \dots + \frac{1}{(2k+1)(2k+3)} = \frac{k}{3(2k+3)} \qquad \dots (1)$$

We shall now prove that P(k + 1) is true.

$$\begin{bmatrix} \frac{1}{3.5} + \frac{1}{5.7} + \frac{1}{7.9} + \dots + \frac{1}{(2k+1)(2k+3)} \end{bmatrix} + \frac{1}{\{2(k+1)+1\}\{2(k+1)+3\}}$$

$$= \frac{k}{3(2k+3)} + \frac{1}{(2k+3)(2k+5)}$$

$$= \frac{1}{(2k+3)} \begin{bmatrix} \frac{k}{3} + \frac{1}{(2k+5)} \end{bmatrix}$$

$$= \frac{1}{(2k+3)} \begin{bmatrix} \frac{k(2k+5)+3}{3(2k+5)} \end{bmatrix}$$

$$= \frac{1}{(2k+3)} \begin{bmatrix} \frac{2k^2+5k+3}{3(2k+5)} \end{bmatrix}$$

$$= \frac{1}{(2k+3)} \begin{bmatrix} \frac{2k^2+2k+3k+3}{3(2k+5)} \end{bmatrix}$$

$$= \frac{1}{(2k+3)} \begin{bmatrix} \frac{2k(k+1)+3(k+1)}{3(2k+5)} \end{bmatrix}$$

$$= \frac{(k+1)(2k+3)}{3(2k+5)}$$

$$= \frac{(k+1)}{3\{2(k+1)+3\}}$$

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

## Q18 :

Prove the following by using the principle of mathematical induction for

all 
$$n \in N$$
:  
 $1+2+3+\ldots+n < \frac{1}{8}(2n+1)^2$ 

#### Answer :

Let the given statement be P(n), i.e.,

$$P(n): 1+2+3+...+n < \frac{1}{8}(2n+1)^2$$

$$e^{1 < \frac{1}{8}(2.1+1)^2 = \frac{9}{8}}$$

It can be noted that P(n) is true for n = 1 since

Let P(k) be true for some positive integer k, i.e.,

$$1+2+\ldots+k < \frac{1}{8}(2k+1)^2 \qquad \ldots (1)$$

We shall now prove that P(k + 1) is true whenever P(k) is true.

Consider

$$(1+2+...+k)+(k+1) < \frac{1}{8}(2k+1)^{2}+(k+1) \qquad [Using(1)]$$

$$<\frac{1}{8}\{(2k+1)^{2}+8(k+1)\}$$

$$<\frac{1}{8}\{4k^{2}+4k+1+8k+8\}$$

$$<\frac{1}{8}\{4k^{2}+12k+9\}$$

$$<\frac{1}{8}\{2k+3)^{2}$$

$$<\frac{1}{8}\{2(k+1)+1\}^{2}$$

 $(1+2+3+...+k)+(k+1)<\frac{1}{8}(2k+1)^2+(k+1)$ 

Hence,

Thus, P(k + 1) is true whenever P(k) is true.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

## Q19:

Prove the following by using the principle of mathematical induction for all  $n \in N$ : n(n + 1)(n + 5) is a multiple of 3.

## Answer :

Let the given statement be P(n), i.e.,

P(n): n(n + 1)(n + 5), which is a multiple of 3.

It can be noted that P(n) is true for n = 1 since 1 (1 + 1) (1 + 5) = 12, which is a multiple of 3.

Let P(k) be true for some positive integer k, i.e.,

k(k+1)(k+5) is a multiple of 3.

:: k (k + 1) (k + 5) = 3m, where  $m \in \mathbb{N} ... (1)$ 

We shall now prove that P(k + 1) is true whenever P(k) is true.

$$(k+1)\{(k+1)+1\}\{(k+1)+5\}$$

$$= (k+1)(k+2)\{(k+5)+1\}$$

$$= (k+1)(k+2)(k+5)+(k+1)(k+2)$$

$$= \{k(k+1)(k+5)+2(k+1)(k+5)\}+(k+1)(k+2)$$

$$= 3m+(k+1)\{2(k+5)+(k+2)\}$$

$$= 3m+(k+1)\{2k+10+k+2\}$$

$$= 3m+(k+1)(3k+12)$$

$$= 3m+(k+1)(k+4)$$

$$= 3\{m+(k+1)(k+4)\} = 3 \times q, \text{ where } q = \{m+(k+1)(k+4)\} \text{ is some natural number Therefore, } (k+1)\{(k+1)+1\}\{(k+1)+5\} \text{ is a multiple of 3.}$$

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

### Q20:

Prove the following by using the principle of mathematical induction for all  $n \in N$ :  $10^{2n-1} + 1$  is divisible by 11.

#### Answer :

Let the given statement be P(n), i.e.,

P(*n*):  $10^{2n} \stackrel{\text{a}e^{-1}}{=} + 1$  is divisible by 11.

It can be observed that P(n) is true for n = 1 since  $P(1) = 10^{2.1 \text{ det } 1} + 1 = 11$ , which is divisible by 11.

Let P(k) be true for some positive integer k, i.e.,

 $10^{2kae^{-1}} + 1$  is divisible by 11.

∴10<sup>2ka€ 1</sup> + 1 = 11*m*, where  $m \in \mathbf{N}$  ... (1)

We shall now prove that P(k + 1) is true whenever P(k) is true.

$$10^{2(k+1)-1} + 1$$
  
=  $10^{2k+2-1} + 1$   
=  $10^{2(k+1)} + 1$   
=  $10^{2} (10^{2k-1} + 1 - 1) + 1$   
=  $10^{2} (10^{2k-1} + 1) - 10^{2} + 1$   
=  $10^{2} \cdot 11m - 100 + 1$  [Using (1)]  
=  $100 \times 11m - 99$   
=  $11(100m - 9)$   
=  $11r$ , where  $r = (100m - 9)$  is some natural number  
Therefore,  $10^{2(k+1)-1} + 1$  is divisible by 11.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

## Q21 :

Prove the following by using the principle of mathematical induction for all  $n \in N$ :  $x^{2n} - y^{2n}$  is divisible by x + y.

#### Answer :

Let the given statement be P(n), i.e.,

P(*n*):  $x^{2n}$  –  $y^{2n}$  is divisible by x + y.

It can be observed that P(n) is true for n = 1.

This is so because  $x^{2 \times 1} \ \hat{a} \in y^{2 \times 1} = x^{2} \ \hat{a} \in y^{2} = (x + y) \ (x \ \hat{a} \in y)$  is divisible by (x + y).

Let P(k) be true for some positive integer k, i.e.,

 $x^{2k}$  â  $\in$  "  $y^{2k}$  is divisible by x + y.

∴ $x^{2k}$  –  $y^{2k} = m (x + y)$ , where  $m \in \mathbb{N}$  ... (1)

We shall now prove that P(k + 1) is true whenever P(k) is true.

$$\begin{aligned} x^{2(k+1)} - y^{2(k+1)} \\ &= x^{2k} \cdot x^2 - y^{2k} \cdot y^2 \\ &= x^2 \left( x^{2k} - y^{2k} + y^{2k} \right) - y^{2k} \cdot y^2 \\ &= x^2 \left\{ m(x+y) + y^{2k} \right\} - y^{2k} \cdot y^2 \qquad \left[ \text{Using (1)} \right] \\ &= m(x+y)x^2 + y^{2k} \cdot x^2 - y^{2k} \cdot y^2 \\ &= m(x+y)x^2 + y^{2k} \left( x^2 - y^2 \right) \\ &= m(x+y)x^2 + y^{2k} \left( x^2 - y^2 \right) \\ &= m(x+y)x^2 + y^{2k} \left( x+y \right) (x-y) \\ &= (x+y) \left\{ mx^2 + y^{2k} \left( x-y \right) \right\}, \text{ which is a factor of } (x+y). \end{aligned}$$

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

## Q22 :

Prove the following by using the principle of mathematical induction for all  $n \in N$ :  $3^{2n+2} - 8n - 9$  is divisible by 8.

## Answer :

Let the given statement be P(n), i.e.,

P(*n*):  $3^{2n+2}$  – 8*n* – 9 is divisible by 8.

It can be observed that P(n) is true for n = 1 since  $3^{2 \times 1+2} \hat{a} \in 0$  × 1  $\hat{a} \in 0$  = 64, which is divisible by 8.

Let P(k) be true for some positive integer k, i.e.,

3<sup>2k+ 2</sup> – 8*k* – 9 is divisible by 8.

∴ $3^{2^{k+2}}$  – 8k – 9 = 8m; where  $m \in \mathbb{N}$  ... (1)

We shall now prove that P(k + 1) is true whenever P(k) is true.

$$3^{2(k+1)+2} - 8(k+1) - 9$$
  
=  $3^{2k+2} \cdot 3^2 - 8k - 8 - 9$   
=  $3^2 (3^{2k+2} - 8k - 9 + 8k + 9) - 8k - 17$   
=  $3^2 (3^{2k+2} - 8k - 9) + 3^2 (8k + 9) - 8k - 17$   
=  $9.8m + 9(8k + 9) - 8k - 17$   
=  $9.8m + 72k + 81 - 8k - 17$   
=  $9.8m + 64k + 64$   
=  $8(9m + 8k + 8)$   
=  $8r$ , where  $r = (9m + 8k + 8)$  is a natural number  
Therefore,  $3^{2(k+1)+2} - 8(k+1) - 9$  is divisible by 8.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

## Q23 :

Prove the following by using the principle of mathematical induction for all  $n \in N$ : 41<sup>*n*</sup> - 14<sup>*n*</sup> is a multiple of 27.

#### Answer :

Let the given statement be P(n), i.e.,

P(*n*):41<sup>*n*</sup> – 14<sup>*n*</sup> is a multiple of 27.

It can be observed that P(n) is true for n = 1 since  $41^{1} - 14^{1} = 27$ , which is a multiple of 27.

Let P(k) be true for some positive integer k, i.e.,

41<sup>k</sup> – 14<sup>k</sup>is a multiple of 27

∴41<sup>*k*</sup> – 14<sup>*k*</sup> = 27*m*, where  $m \in \mathbf{N}$  ... (1)

We shall now prove that P(k + 1) is true whenever P(k) is true.

$$41^{k+1} - 14^{k+1}$$
  
=  $41^{k} \cdot 41 - 14^{k} \cdot 14$   
=  $41(41^{k} - 14^{k} + 14^{k}) - 14^{k} \cdot 14$   
=  $41(41^{k} - 14^{k}) + 41.14^{k} - 14^{k} \cdot 14$   
=  $41.27m + 14^{k} (41 - 14)$   
=  $41.27m + 27.14^{k}$   
=  $27(41m - 14^{k})$   
=  $27 \times r$ , where  $r = (41m - 14^{k})$  is a natural number  
Therefore,  $41^{k+1} - 14^{k+1}$  is a multiple of 27.

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.

## Q24 :

Prove the following by using the principle of mathematical induction for all  $n \in \mathbb{N}$  :

 $(2n+7) < (n+3)^2$ 

### Answer :

Let the given statement be P(n), i.e.,

$$P(n): (2n+7) < (n+3)^2$$

It can be observed that P(n) is true for n = 1 since  $2 \cdot 1 + 7 = 9 < (1 + 3)^2 = 16$ , which is true.

Let P(k) be true for some positive integer k, i.e.,

 $(2k+7) < (k+3)^2 \dots (1)$ 

We shall now prove that P(k + 1) is true whenever P(k) is true.

$$\{2(k+1)+7\} = (2k+7)+2 
\therefore \{2(k+1)+7\} = (2k+7)+2 < (k+3)^2 + 2 \qquad [u \sin g (1)] 
2(k+1)+7 < k^2 + 6k + 9 + 2 
2(k+1)+7 < k^2 + 6k + 11 
Now, k^2 + 6k + 11 < k^2 + 8k + 16 
\therefore 2(k+1)+7 < (k+4)^2 
2(k+1)+7 < {(k+1)+3}^2$$

Hence, by the principle of mathematical induction, statement P(n) is true for all natural numbers i.e., n.